

**If you find you are taking large amounts of pain medicine at frequent intervals, please call our office.** If you anticipate needing more prescription medication for the weekend, we appreciate your making every effort to call for a refill during weekday business hours. In the unlikely situation that you need a prescription over the weekend, please call our emergency number.

**NAUSEA:** Nausea is not uncommon after an extraction. Sometimes pain medications are the cause. Nausea can be reduced by preceding each pain pill with a small amount of soft food, and taking the pill with a large volume of water. Try to keep taking clear fluids and minimize dosing of pain medications, but call us if you do not feel better. Classic Coca Cola may help with nausea.

**DIET:** Eat any nourishing food that can be taken with comfort. Avoid extremely hot foods. Do not use a straw for the first few days after surgery. It is sometimes advisable, but not absolutely required, to confine the first day's intake to liquids or pureed foods (soups, puddings, yogurt, milk shakes, etc.) It is best to avoid foods like nuts, sunflower seeds, popcorn, etc., which may get lodged in the socket areas. Over the next several days you may gradually progress to solid foods. It is important not to skip meals! If you take nourishment regularly you will feel better, gain strength, have less discomfort and heal faster. If you are a diabetic, maintain your normal eating habits or follow instructions given by your doctor.

**SHARP EDGES:** If you feel something hard or sharp edges in the extraction areas, it is likely you are feeling the bony walls which once supported the extracted teeth. Occasionally small slivers of bone may work themselves out during the following week or so. If they cause concern or discomfort, please call the office.

#### **INSTRUCTIONS FOR THE SECOND AND THIRD DAYS**

**MOUTH RINSES:** Keeping your mouth clean after extractions is essential. Use 1/4 teaspoon of salt dissolved in an 8 ounce glass of warm water and gently rinse with portions of the solution, taking five minutes to use the entire glassful. Repeat as often as you like, but at least two or three times daily.

**BRUSHING:** Begin your normal oral hygiene routine as soon as possible after the procedure. Soreness and swelling may not permit vigorous brushing, but please make every effort to clean your teeth within the bounds of comfort.

**HOT APPLICATIONS:** You may apply warm compresses to the skin over the areas of swelling (hot water bottle, hot moist towels, heating pad) for 20 minutes on and 20 minutes off to help soothe tender areas. This will also help decrease swelling and stiffness.

**HEALING:** Normal healing after tooth extraction should be as follows: The first two days after surgery are generally the most uncomfortable and there is usually some swelling. On the third day you should be more comfortable and, although still swollen, can usually begin a more substantial diet. **The remainder of the post-procedure course should be gradual, steady improvement.** If you don't see continued improvement, please call our office. If you are given a plastic irrigating syringe, **DO NOT** use it for the first five days. Then use it daily according to the instructions until you are certain the tooth socket has closed completely and that there is no chance of any food particles lodging in the socket.

It is our desire that your recovery be as smooth and pleasant as possible. Following these instructions will assist you, but if you have questions about your progress, please call the office where you had your procedure. A 24-hour answering service is available to contact the doctor on call after hours. Calling during office hours will afford a faster response to your question or concern.